

Asymptotics of Elliptic Genera of Symmetric Products and Dyonic Black Hole Degeneracy

A.A. Bytsenko ^{*}, A.E. Gonçalves [†]

Departamento de Fisica, Universidade Estadual de Londrina,
Caixa Postal 6001, Londrina-Parana, Brazil

and

S.D. Odintsov [‡]

Departamento de Fisica, Universidad del Valle,
AA 25360, Cali, Colombia

February 7, 2008

Abstract

We calculate asymptotic expansions of elliptic genera for a supersymmetric sigma model on the N -fold symmetric product $S^N M$ of a Kähler manifold M and for $N = 2$ superconformal field theory. Asymptotic expansions for the degeneracy of dyonic black hole spectrum are also derived.

I. In this note we use an identity proved in [1] that equates elliptic genus partition function of a supersymmetric sigma model on the N -fold symmetric product $S^N M$ of M ($S^N M = M^N/S_N$, S_N is the symmetric group of N elements) to the partition function of a second quantized string

^{*}E-mail: abyts@fisica.uel.br On leave from Sankt-Petersburg State Technical University, Russia

[†]E-mail: goncalve@fisica.uel.br

[‡]E-mail: odintsov@quantum.univalle.edu.co On leave from Tomsk Pedagogical University, Russia

theory on the space $S^1 \otimes M$. The generating function of these elliptic genera is almost an automorphic form for $O(3, 2, \mathbb{Z})$ T -duality group. We shall derive asymptotic expansion of elliptic genus as well as asymptotic growth for the degeneracy of dyonic spectrum in four-dimensional $N = 4$ string theory. It is known that the counting formula is manifestly symmetric under the duality group; we wish to present its asymptotic growth in explicit form, which reproduces the macroscopic Bekenstein-Hawking entropy.

We start with a supersymmetric sigma model on $S^N M$. The Hilbert space of an orbifold field theory can be decomposed into twisted sectors $\mathcal{H}_\gamma$, that are labelled by the conjugacy classes $\{\gamma\}$ of the orbifold group S_N [2, 3, 1]. For given twisted sector one can keep the states invariant under the centralizer subgroup Γ_γ related to the element γ . Let $\mathcal{H}_\gamma^{\Gamma_\gamma}$ is an invariant subspace associated with Γ_γ ; the total orbifold Hilbert space takes the form $\mathcal{H}(S^N M) = \bigoplus_{\{\gamma\}} \mathcal{H}_\gamma^{\Gamma_\gamma}$. Taking into account the group S_N one can compute the conjugacy classes $\{\gamma\}$ by set of partitions $\{N_n\}$ of N , namely $\sum_n n N_n = N$, where N_n is the multiplicity of the cyclic permutation (n) of n elements in the decomposition of γ : $\{\gamma\} = \sum_{j=1}^s (j)^{N_j}$. For this conjugacy class the centralizer subgroup of a permutation γ is $\Gamma_\gamma = S_{N_1} \otimes_{j=2}^s (S_{N_j} \rtimes \mathbb{Z}_j^{N_j})$ [1], where each subfactor S_{N_n} and $\mathbb{Z}_n$ permutes the N_n cycles (n) and acts within one cycle (n) correspondingly. Following the lines of Ref. [1] we may decompose each twisted sector $\mathcal{H}_\gamma^{\Gamma_\gamma}$ into the product over the subfactors (n) of N_n -fold symmetric tensor products, $\mathcal{H}_\gamma^{\Gamma_\gamma} = \bigotimes_{n>0} S^{N_n} \mathcal{H}_{(n)}^{\mathbb{Z}_n}$, where $S^N \mathcal{H} \equiv (\bigotimes^N \mathcal{H})^{S_N}$. Let $\chi(\mathcal{H}; q, y)$ be the partition function for every (sub) Hilbert space of a supersymmetric sigma model; $q = e[\tau]$, $y = e[z]$, and $e[x] \equiv \exp[2\pi i x]$. It can be shown that the partition function coincides with the elliptic genus [4, 5, 6, 7, 8, 9].

In general the genus is a homomorphism: $\Omega \mapsto \mathbb{C}$, where Ω is the Thom ring of oriented cobordisms. The following relations for the genus hold:

$$\chi(M_1 \sqcup M_2; q, y) = \chi(M_1; q, y) + \chi(M_2; q, y), \quad (1)$$

$$\chi(M_1 \otimes M_2; q, y) = \chi(M_1; q, y) \cdot \chi(M_2; q, y), \quad (2)$$

$$\chi(M_1; q, y) = \chi(M_2; q, y) \text{ if } M_1 \text{ and } M_2 \text{ are cobordant.} \quad (3)$$

The elliptic genus admits the following identities:

$$\chi(\mathcal{H} \oplus \mathcal{H}'; q, y) = \chi(\mathcal{H}; q, y) + \chi(\mathcal{H}'; q, y), \quad (4)$$

$$\chi(\mathcal{H} \otimes \mathcal{H}'; q, y) = \chi(\mathcal{H}; q, y) \cdot \chi(\mathcal{H}'; q, y). \quad (5)$$

For any vector bundle V over a smooth manifold we can define formal sums

$$\bigwedge_q V = \bigoplus_{k \geq 0}^d q^k \bigwedge^k V, \quad S_q V = \bigoplus_{k \geq 0}^{\infty} q^k S^k V, \quad \chi_y(T_M) = \bigoplus_{s=0}^d (-y)^s \chi \left(\bigwedge^s T_M \right), \quad (6)$$

where T_M is the holomorphic tangent bundle of Kähler manifold M of complex dimension d , $\bigwedge^k$ (S^k) denote the k -th exterior (symmetric product). Note that for a Calabi-Yau space the χ_y -genus [10] is a weak Jacobi form of weight zero and index $d/2$ and it transforms as $\chi_y(T_M) = (-1)^{r-d} y^r \chi_{y^{-1}}(T_M)$. This relation can also be derived from the Serre duality $H^j(M; \bigwedge^s T_M) \cong H^{d-j}(M; \bigwedge^{r-s} T_M)$.

The Riemann-Roch-Hirzebruch theorem:

$$\chi(M; q, y) \stackrel{def}{=} \sum (-1)^j \dim H^j(M; E_{q,y}) = \int_M \text{ch}(E_{q,y}) Td(M), \quad (7)$$

gives an equivalent definition of the elliptic genus. The bundle $E_{q,y}$ of a characteristic class is

$$E_{q,y} = y^{-\frac{d}{2}} \bigotimes_{n \geq 1}^{\infty} \left(\bigwedge_{-yq^{n-1}} T_M \bigotimes_{n \geq 1}^{\infty} \bigwedge_{-y^{-1}q^n} T_M^* \bigotimes_{n \geq 1}^{\infty} S_{q^n} T_M \bigotimes_{n \geq 1}^{\infty} S_{q^n} T_M^* \right), \quad (8)$$

and the integral (7) can be expanded as

$$\begin{aligned} \chi(M; q, y) &= \chi_y(T_M) + q \left\{ \bigoplus_{s=0}^d [(-y)^{s+1} \chi \left(\bigwedge^s T_M \otimes T_M \right) \right. \\ &\quad \left. + (-y)^{s-1} \chi \left(\bigwedge^s T_M \otimes T_M^* \right) + (-y)^s \chi \left(\bigwedge^s T_M \otimes (T_M \otimes T_M^*) \right)] \right\} + \mathcal{O}(q^{3/2}), \end{aligned} \quad (9)$$

where we used formula (7). Note that the elliptic genus was independently introduced in [11, 12]. For $q = 0$ the elliptic genus reduces to a weighted sum over the Hodge numbers, namely $\chi(M; 0, y) = \sum_{p,q} (-1)^{p+q} y^{p-\frac{d}{2}} h^{p,q}(M)$.

If $\chi(\mathcal{H}_{(n)}^{\mathbb{Z}_n}; q, y)$ admits the extension $\chi(\mathcal{H}; q, y) = \sum_{m \geq 0, \ell} C(nm, \ell) q^m y^\ell$, the following result holds (see Ref. [1, 13]):

$$\sum_{N \geq 0} p^N \chi(S^N \mathcal{H}_{(n)}^{\mathbb{Z}_n}; q, y) = \prod_{m \geq 0, \ell} (1 - pq^m y^\ell)^{-C(nm, \ell)}, \quad (10)$$

$$\sum_{N \geq 0} p^N \chi(S^N M; q, y) = \prod_{n > 0, m \geq 0, \ell} (1 - p^n q^m y^\ell)^{-C(nm, \ell)}. \quad (11)$$

Here $p = \mathbf{e}[\rho]$, ρ and τ determine the complex Kähler form and complex structure modulus of T^2 respectively, and z parametrizes the $U(1)$ bundle on T^2 . The logarithm of the partition function $Z(p; q, y)$ is the one-loop free energy $F(p; q, y)$ for a string on $T^2 \otimes M$:

$$\begin{aligned} F(p; q, y) &= \log Z(p; q, y) = - \sum_{n > 0, m, \ell} C(nm, \ell) \log(1 - p^n q^m y^\ell) \\ &= \sum_{n > 0, m, \ell, k > 0} \frac{1}{k} C(nm, \ell) p^{kn} q^{km} y^{k\ell} = \sum_{N > 0} p^N \sum_{kn=N} \frac{1}{k} \sum_{m, \ell} C(nm, \ell) q^{km} y^{k\ell}. \end{aligned} \quad (12)$$

The free energy can be written as a sum of Hecke operators T_N [14] acting on the elliptic genus of M [1, 15, 16]: $F(p; q, y) = \sum_{N > 0} p^N T_N \chi(M; q, y)$.

II. The goal now is to calculate an asymptotic expansion of the elliptic genus $\chi(S^N M; q, y)$. The Laurent inversion formula has the form

$$\chi(S^N M; q, y) = \frac{1}{2\pi i} \oint \frac{Z(p, q, y)}{p^{N+1}} dp, \quad (13)$$

where the contour integral is taken on a small circle about the origin. Let the Dirichlet series

$$\mathfrak{D}(s; \tau, z) = \sum_{(n, m, \ell) > 0} \sum_{k=1}^{\infty} \frac{\mathbf{e}[\tau m k + z \ell k] C(nm, \ell)}{n^s k^{s+1}} \quad (14)$$

converges for $0 < \Re s < \alpha$. We assume that series (14) can be analytically continued in the region $\Re s \geq -C_0$ ($0 < C_0 < 1$) where it is analytic except a pole of order one at $s = 0$ and $s = \alpha$ with residue $\text{Res}[\mathfrak{D}(0; \tau, z)]$ and $\text{Res}[\mathfrak{D}(\alpha; \tau, z)]$ respectively. Besides let $\mathfrak{D}(s; \tau, z) = \mathcal{O}(|\Im s|^{C_1})$ uniformly in $\Re s \geq -C_0$ as $|\Im s| \rightarrow \infty$, where C_1 is a fixed positive real number. If $t \equiv 2\pi(\Im \rho - i \Re \rho)$ then the Mellin-Barnes representation of the function $F(t; \tau, z)$ gives

$$\mathfrak{M}[F](t; \tau, z) = \frac{1}{2\pi i} \int_{\Re s = 1 + \alpha} t^{-s} \Gamma(s) \mathfrak{D}(s; \tau, z) ds. \quad (15)$$

The integrand in Eq. (17) has a first order pole at $s = \alpha$ and a second order pole at $s = 0$. Shifting the vertical contour from $\Re s = 1 + \alpha$ to $\Re s = -C_0$

(this procedure is permissible) and making use of the residues theorem one obtains

$$F(t; \tau, z) = t^{-\alpha} \Gamma(\alpha) \text{Res}[\mathfrak{D}(\alpha; \tau, z)] + \lim_{s \rightarrow 0} \frac{d}{ds} [s \mathfrak{D}(s; \tau, z)] - (\gamma + \log t) \text{Res}[\mathfrak{D}(0; \tau, z)] + \frac{1}{2\pi i} \int_{\Re s = -C_0} t^{-s} \Gamma(s) \mathfrak{D}(s; \tau, z) ds. \quad (16)$$

The absolute value of the integral in (16) can be estimated to behave as $\mathcal{O}((2\pi \Im \rho)^{C_0})$.

We are ready now to state the result, which follows in a similar way from the Meinardus main theorem [17, 18, 19]. In the half-plane $\Re t > 0$ there exists an asymptotic expansion for $Z(t; \tau, z)$ uniformly in $|\Re \rho|$ for $|\Im \rho| \rightarrow 0$, $|\arg(2\pi i \rho)| \leq \pi/4$, $|\Re \rho| \leq 1/2$ and given by

$$Z(t; \tau, z) = e^{\left[\frac{1}{2\pi i} \left\{ \text{Res}[\mathfrak{D}(\alpha; \tau, z)] \Gamma(\alpha) t^{-\alpha} - \text{Res}[\mathfrak{D}(0; \tau, z)] \log t - \gamma \text{Res}[\mathfrak{D}(0; \tau, z)] + \lim_{s \rightarrow 0} \frac{d}{ds} [s \mathfrak{D}(s; \tau, z)] + \mathcal{O}(|2\pi \Im \tau|^{C_0}) \right\} \right]}. \quad (17)$$

The asymptotic expansion for the elliptic genus has the form

$$\chi(S^N M; \tau, z)_{N \rightarrow \infty} = \mathcal{C}(\alpha; \tau, z) N^{(2\text{Res}[\mathfrak{D}(0; \tau, z)] - 2 - \alpha)/(2(1 + \alpha))} \times e^{\left[\frac{1 + \alpha}{2\pi i \alpha} (\text{Res}[\mathfrak{D}(\alpha; \tau, z)] \Gamma(1 + \alpha))^{1/(1 + \alpha)} N^{\alpha/(1 + \alpha)} \right]} [1 + \mathcal{O}(N^{-k})], \quad (18)$$

$$\mathcal{C}(\alpha; \tau, z) = \{\text{Res}[\mathfrak{D}(\alpha; \tau, z)] \Gamma(1 + \alpha)\}^{(1 - 2\text{Res}[\mathfrak{D}(0; q, y)])/(2 + 2\alpha)} \times e^{\left[\frac{1}{2\pi i} \left(\lim_{s \rightarrow 0} \frac{d}{ds} [s \mathfrak{D}(0; \tau, z)] - \gamma \text{Res}[\mathfrak{D}(0; \tau, z)] \right) \right]} [2\pi(1 + \alpha)]^{1/2}, \quad (19)$$

where k is a positive constant (see for detail Refs. [19, 20, 21]). In above formulae the complete form of the prefactor $\mathcal{C}(\alpha; \tau, z)$ appears. The result (18), (19) has an universal character for all elliptic genera associated to Calabi-Yau manifolds.

Finally we go into some facts related to orbifoldized elliptic genus of $N = 2$ superconformal field theory. The contribution of the untwisted sector to the orbifoldized elliptic genus is the function $\chi(M; \tau, z) \equiv \phi(\tau, z) \equiv$ $\begin{array}{c} \square \\ 0 \end{array}(\tau, z)$,

whereas

$$\phi\left(\frac{a\tau+b}{c\tau+d}, \frac{z}{c\tau+d}\right) = {}_0 \square_0(\tau, z) \mathbf{e}\left[\frac{rcz^2}{c\tau+d}\right], \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}), \quad (20)$$

$r = d/2$. The contribution of the twisted μ - sector projected by ν is [9]:

$${}_\nu \square_\mu(\tau, z) = \phi(\tau, z + \mu\tau + \nu) \mathbf{e}\left[\frac{d}{2}(\mu\nu + \mu^2\tau + 2\mu z)\right], \quad \mu, \nu \in \mathbb{Z}. \quad (21)$$

The orbifoldized elliptic genus can be defined by

$$\phi(\tau, z)_{\text{orb}} \stackrel{\text{def}}{=} \frac{1}{h} \sum_{\mu, \nu=0}^{h-1} (-1)^{P(\mu+\nu+\mu\nu)} {}_\nu \square_\mu(\tau, z), \quad (22)$$

where P, h are some integers. Using transformation properties (21) in Eqs. (18), (19) one can obtain the asymptotic expansion for the orbifoldized elliptic genus. In fact we can introduce a procedure, starting with the expansion of the elliptic genus of the untwisted sector, to compute the asymptotics of the elliptic genus of the twisted sector.

III. Let T and U are the Kähler and complex structure moduli of the torus $\mathbf{T}^2$. The Narain duality group $SO(2, 3; \mathbb{Z})$ is isomorphic to Siegel modular group $Sp(4, \mathbb{Z})$ and it is convenient to combine the parameters T, U and a Wilson line modules V into $2 \otimes 2$ matrix belonging to Siegel upper half plane of genus 2

$$\Omega = \begin{pmatrix} T & V \\ V & U \end{pmatrix}, \quad \Im T, \Im U > 0, \quad \det(\Im \Omega) > 0. \quad (23)$$

One can use Θ -transform [22, 23], which converts the $K3$ elliptic genus $\chi(K3; q, y) = \sum_{m, \ell} C(4m - \ell^2) \mathbf{e}[m\tau + \ell z]$ to the automorphic form [16, 24, 25, 26, 27]

$$\Phi(T, U, V) = \mathbf{e}[T + U + V] \prod_{(n, m, \ell) > 0} (1 - \mathbf{e}[nT + mU + \ell V])^{C(4nm - \ell^2)}. \quad (24)$$

This form is the denominator product for the GKM based on the Cartan matrix

$$\mathcal{A} = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}. \quad (25)$$

A counting formula for the degeneracies of a black hole preserving 1/4 of the supersymmetry of the background $IIA/K3 \otimes T^2$, or heterotic/ T^6 is presented in [13]. The degeneracies for the supersymmetric black holes with (electric, magnetic) charge vectors $\mathbf{q} = (q_e, q_m) \in II^{6,22} \oplus II^{6,22}$ have the form

$$d(q_e, q_m) = \oint \frac{\mathbf{e} \left[\frac{1}{2} \mathbf{q} \cdot \Omega \cdot \mathbf{q}^t \right]}{\Phi(T, U, V)} dT dU dV. \quad (26)$$

Let us expand the integrand of Eq. (26) as a formal Fourier series expansion

$$\mathbf{e} \left[\frac{\mathcal{F}(\Omega)}{2\pi i} \right] \equiv \frac{\mathbf{e}[T + U + V]}{\Phi(\Omega)} = \sum_{(n, m, \ell)} \mathbf{D}(n, m, \ell) \mathbf{e}[-(nT + mU + \ell V)], \quad (27)$$

where (n, m, ℓ) are integers. Let

$$\begin{aligned} \mathfrak{D}_1(s; n, m) &= \sum_{\ell} \frac{C(nm, \ell)}{\ell^s}, \quad \mathfrak{D}_1(s; n) = \sum_m \frac{\text{Res}[\mathfrak{D}_1(\alpha_3; n, m)]}{m^s}, \\ \mathfrak{D}_1(s) &= \sum_n \frac{\text{Res}[\mathfrak{D}_1(\alpha_2; n)]}{n^s} \end{aligned} \quad (28)$$

be the Dirichlet series which converges for $0 < \Re s < \alpha_j$ ($j = 1, 2, 3$). We assume that the series $\mathfrak{D}_1(s; n, m)$, $\mathfrak{D}_1(s; n)$ and $\mathfrak{D}_1(s)$ have poles of order one at $s = \alpha_3, \alpha_2$ and α_1 respectively. We assume also that for $r \in \mathbb{Z}_+$ the series

$$\begin{aligned} \mathfrak{D}_2(s; -r) &= \sum_n \frac{\mathfrak{D}_1(-r; n)}{n^s}, \quad \mathfrak{D}_2(s; n, -r) = \sum_m \frac{\mathfrak{D}_1(-r; n, m)}{m^s}, \\ \mathfrak{D}_2(s; \beta_2, -r) &= \sum_n \frac{\text{Res}[\mathfrak{D}_2(\beta_2; n, -r)]}{n^s} \end{aligned} \quad (29)$$

converges for $0 < \Re s < \beta_j$ and have poles of order one at $s = \beta_1, \beta_2$ and β_3 respectively.

If $\mathcal{F}(\Omega)$ satisfy the expansion (27) and $\mathbf{e}[-(T+U+V)] \equiv \exp[W+X+Y]$ then it can be shown that

$$\mathbf{e}\left[\frac{\mathcal{F}(\Omega)}{2\pi i}\right] = \text{Res}[\mathfrak{D}_1(\alpha_1)] \prod_{j=1}^3 \Gamma(\alpha_j) W^{-\alpha_1} X^{-\alpha_2} Y^{-\alpha_3} \zeta_R(\sigma) \\ + \mathcal{O}\left(\max\left\{X^{-\alpha_2} Y^{\alpha_3}, W^{-\beta_1} Y^{-\alpha_3}, W^{-\beta_3} X^{-\beta_2}\right\}\right), \quad (30)$$

uniformly in $\{\Im W, \Im X, \Im Y\}$ as $\{\Re W, \Re X, \Re Y\} \rightarrow 0$. Here $\zeta_R(\sigma)$ is the Riemann zeta function, $\sigma = 1 + \alpha_1 + \alpha_2 + \alpha_3$. For $\{n, m, \ell\} \rightarrow \infty$ one has

$$\mathbf{D}(n, m, \ell) \simeq \mathcal{I}(n, m, \ell) \mathbf{e} \left[\frac{\sigma - 1}{2\pi i} \left(\text{Res}[\mathfrak{D}_1(\alpha_1)] \prod_{j=1}^3 \Gamma(\alpha_j) \zeta_R(\sigma) \right. \right. \\ \left. \left. \times \left(\frac{n}{\alpha_1}\right)^{\alpha_1} \left(\frac{m}{\alpha_2}\right)^{\alpha_2} \left(\frac{\ell}{\alpha_3}\right)^{\alpha_3} \right)^{\frac{1}{\sigma}} \right], \quad (31)$$

and

$$\mathcal{I}(n, m, \ell) = \int_{\mathcal{G}} \mathbf{e} \left[\frac{\mathcal{F}(\Omega) + i(n\Im W + m\Im X + \ell\Im Y)}{2\pi i} \right] \frac{d(\Im W) d(\Im X) d(\Im Y)}{(2\pi)^3}, \quad (32)$$

$$\mathcal{G} = \{|\Re W|, |\Re X|, |\Re Y| < \epsilon < 1\}. \quad (33)$$

Using Eqs. (31) and (32) one can obtain the degeneracy for a given electric and magnetic charge. The Eq. (31) gives an explicit counting formula of the asymptotic of microscopic black hole states degeneracy and in particular allows one to analyze the asymptotic growth in more directions. This result is similar to the correct asymptotic growth as predicted from the macroscopic entropy formulas of extremal dyonic black holes.

There are intriguing relations between string duality, the theory of generalized Kac-Moody algebras, and their associated automorphic forms. For example, threshold corrections involve automorphic forms similar to those associated to Kac-Moody algebras; black hole BPS states are counted by a denominator product for a Kac-Moody algebra. In this note we used some of these relations in order to give the correct asymptotic expansion for the degeneracies. We hope that proposed calculation will be interesting in view of future applications to concrete problems in string theory and in black hole physics.

Acknowledgements

A.A. Bytsenko wishes to thank CNPq and the Department of Physics of Londrina University for financial support and kind hospitality. The research of A.A. Bytsenko was supported in part by Russian Foundation for Basic Research (grant No. 98-02-18380-a) and by GRACENAS (grant No. 6-18-1997). The work of S.D. Odintsov has been supported in part by Universidad del Valle (Colombia) and GRACENAS (grant No. 6-18-1997).

References

- [1] R. Dijkgraaf, G. Moore, E. Verlinde and H. Verlinde, Commun. Math. Phys. **185**, 197 (1997).
- [2] L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl. Phys. **B 261**, 620 (1985).
- [3] L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl. Phys. **B 274**, 285 (1986).
- [4] A. Schellekens and N. Warner, Phys. Lett. **B 177**, 317 (1986).
- [5] A. Schellekens and N. Warner, Nucl Phys. **B 287**, 317 (1987).
- [6] E. Witten, Commun. Math. Phys. **109**, 525 (1987).
- [7] P.S. Landweber Ed., “*Elliptic Curves and Modular Forms in Algebraic Topology*”, Springer-Verlag (1988).
- [8] T. Eguchi, H. Ooguri, A. Taormina and S.-K. Yang, Nucl. Phys. **B 315**, 193 (1989).
- [9] T. Kawai, Y. Yamada and S.-K. Yang, Nucl. Phys. **B 414**, 191 (1994).
- [10] F. Hirzebruch, “*Topological Methods in Algebraic Geometry*”, 3rd Edition, Springer-Verlag (1978).
- [11] O. Alvarez, T.P. Killingback, M. Mangano and P. Windey, Commun. Math. Phys. **111**, 1 (1987).

- [12] O. Alvarez, T.P. Killingback, M. Mangano and P. Windey, “*The Dirac-Ramond Operator in String Theory and Loop Space Index Theorems*”, Nucl. Phys. **B** (Proc. Suppl.) **1A** (1988).
- [13] R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. **B** **484**, 543 (1997).
- [14] S. Lang, “*Introduction to Modular Forms*”, Springer-Verlag (1976).
- [15] R.E. Borcherds, Invent. Math. **120**, 161 (1995).
- [16] V.A. Gritsenko and V.V. Nikulin, “*Siegel Automorphic Form Corrections of Some Lorentzian Kac-Moody Algebras*”, alg-geom/9504006 (1995).
- [17] G. Meinardus, Math. Z. **59**, 338 (1954).
- [18] G. Meinardus, Math. Z. **61**, 289 (1954).
- [19] G.E. Andrews, “*The Theory of Partitions*”. In Encyclopedia of Mathematics and its Applications, Addison-Wesley Publishing Company (1976).
- [20] E. Elizalde, S.D. Odintsov, A. Romeo, A.A. Bytsenko and S. Zerbini, “*Zeta Regularization Techniques with Applications*”, World Scientific, Singapore (1994).
- [21] A.A. Bytsenko, G. Cognola, L. Vanzo and S. Zerbini, Phys. Rep. **266**, 1 (1996).
- [22] T. Kawai, “*N=2 Heterotic String Threshold Correction, K3 Surface and Generalized Kac-Moody Superalgebra*”, hep-th/9512046 (1995).
- [23] C.D.D. Neumann, “*The Elliptic Genus of Calabi-Yau 3- and 4-Folds, Product Formulae and Generalized Kac-Moody Algebras*”, hep-th/9607029 (1996).
- [24] V.A. Gritsenko and V.V. Nikulin, “*K3 Surfaces, Lorentzian Kac-Moody Algebras, and Mirror Symmetry*”, alg-geom/9510008 (1995).

- [25] V.A. Gritsenko and V.V. Nikulin, “*Automorphic Forms and Lorentzian Kac-Moody Algebras. Part I*”, alg-geom/9610022 (1996).
- [26] V.A. Gritsenko and V.V. Nikulin, “*Automorphic Forms and Lorentzian Kac-Moody Algebras. Part II*”, alg-geom/9611028 (1996).
- [27] V.A. Gritsenko and V.V. Nikulin, “*A Lecture on Arithmetic Mirror Symmetry and Calabi-Yau Manifolds*”, alg-geom/9612002 (1996).